

1E2401

1E2401

B.Tech. I Semester (Main) Examination, Dec. - 2018

BSC

1FY2-01 Engineering Mathematics - I

Time : 3 Hours

Maximum Marks : 160

Instructions to Candidates:

Attempt all **ten** questions from **Part A**, any **five** questions out of seven from **Part B** and any **four** questions out of five from **Part C**. (Schematic diagrams must be shown wherever necessary). Any data you feel missing suitably be assumed and stated clearly. Units of quantities used/calculated must be stated clearly.

Part - A

(Answer should be given up to 25 words only). All questions are compulsory.

(10×3=30)

1. What is the value of $\Gamma\left(-\frac{1}{2}\right)$.
2. Find the value of $\int_0^{\pi/2} \sin^6 \theta \cos^7 \theta d\theta$.
3. Find whether series $\sum \frac{n}{n+10}$ is convergent or not?
4. Give an example of two divergent series whose sum is convergent.
5. Find sum of Fourier series of $f(x)$ at $x = 2$ where $f(x) = \begin{cases} 0, & 0 \leq x < 1 \\ 1, & 1 \leq x < 2 \end{cases}$.
6. State Parseval's Theorem.
7. Give an example of two variable function whose both partial derivatives exist but limit does not exist at origin.

8. Find the directions in which $f(x,y) = \frac{x^2}{2} + \frac{y^2}{2}$ increases most rapidly at the point (1,1).

9. Suppose the force field $F = \nabla f$ is the gradient of the function $f(x,y,z) = -\frac{1}{(x^2 + y^2 + z^2)}$. Find the work done by F in moving an object along a smooth curve C joining (1,0,0) to (0,0,2) that does not pass through origin.

10. Find $\iint_S \vec{r} \cdot \hat{n} \, dS$ where S is a closed surface enclosing volume V and $\vec{r} = xi + yj + zk$.

Part - B

(Analytical/Problem solving questions). Attempt any five questions.

(5×10=50)

- Find volume of the solid generated by the revolution of the curve $x = a \cos^3 \theta, y = a \sin^3 \theta$ about the x-axis.
- Find Taylor series expansion of $f(x) = \cos 5x^2$ about the point $x = \pi$.
- Obtain half range sine series for $f(x) = e^x$, in $0 < x < 1$.
- If resistors of R_1, R_2 and R_3 ohms are connected in parallel to make an R-ohm resistor, find the value of $\partial R / \partial R_2$ when $R_1 = 30, R_2 = 45$ and $R_3 = 90$ ohms.
- The derivative of $f(x,y)$ at $P_0(1,2)$ in the direction of $i+j$ is $2\sqrt{2}$ and in the direction of $-2j$ is -3 . What is the derivative of f in the direction of $-i-2j$?
- Find the area of the region R in the xy-plane enclosed by the circle $x^2+y^2=4$ above the line $y = 1$ and below the line $y = \sqrt{3}x$.
- Find the centroid of the region in the first quadrant that is bounded above by the line $y = x$ and below by the parabola $y = x^2$.

Part - C

(Descriptive/Analytical/Problem Solving/Design question). Attempt any four questions.

(4×20=80)

- Find the value of $\int_0^\infty \cos x^2 \, dx$.

2. Discuss the convergence of the series $\sum \frac{\sqrt{n}}{\sqrt{n^2+1}} x^n$.

3. Find Fourier series representation of $f(x) = \begin{cases} 0, & -\pi \leq x \leq 0 \\ \sin x, & 0 < x < \pi \end{cases}$ and prove that

$$\frac{1}{1.3} + \frac{1}{3.5} + \frac{1}{5.7} + \dots = \frac{1}{2}$$

4. The plane $x + y + z = 1$ cuts the cylinder $x^2 + y^2 = 1$ in an ellipse. Find the points on the ellipse that lie closest to and farthest from origin.

5. Verify Stoke's theorem for the hemisphere $S: x^2 + y^2 + z^2 = 9, z \geq 0$, its bounding circle $C: x^2 + y^2 = 9, z = 0$, and the field $\vec{F} = y\vec{i} - x\vec{j}$.

